

Ivermectin 12mg Introduction

[Ivermectin 12mg](#) medication is used to treat certain parasitic roundworm diseases like intestinal strongyloidiasis (infection in the gut caused by a type of roundworm called *Strongyloides stercoralis*). An infection caused by *Wuchereria bancrofti* in blood and tissues and human scabies (an infectious disease of the skin) after previous treatment has failed. . Your quality of life can be improved by curing parasite illnesses. Ivermectin tablet contains Iverheal which belongs to the group of medicines called Antiparasitic agents. Ivermectin 12mg tablets are also used for the treatment of river blindness, a parasitic disease caused by *Onchocerca volvulus*(Onchocerciasis).



This Ivermectin 12mg mainly works to kill the parasitic worms in the body. Before taking Ivermectin 12mg tablet, always be concerned with your doctor. Because if you have an eye infection or a weak immune system, you have to face some side effects. Also if you take this medicine with any other drugs for other conditions then this medication does not work.

Ivermectin 12mg tablet is not used for children weighing less than 15kg. In this case, medicine did not work properly. Also did not get the safety and effectiveness of the

Ivermectin 12mg tablet. Because Ivermectin 12mg tablet is mainly used for elderly patients (65 years and above). For more information, you can follow us on [Twitter!](#)

Ivermectin 12mg Side Effects

Common side effects:

- Mild pruritus (itchy skin)
- Feel giddiness
- Vomit sensation
- Pain in abdomen
- Difficulty in moving
- Muscle pain or stiffness
- Difficulty in passing stools
- Lethargy (feeling low or lack of energy)
- Transient changes in electroencephalography (ECG)

Serious side effects:

- Fever
- Hepatitis
- Neurotoxicity
- Loss of vision
- Unusual sleepiness
- Decreased heart rate
- Increase asthma condition
- Unusual bleeding or bruising
- Unusual tiredness or weakness
- Unusual weight gain or loss
- Bleeding of the eye membrane
- Uncontrolled electrical disturbance in the brain

Rare side effects:

- Yellow eyes and skin
- Diarrhea
- Tearing
- Loss of consciousness
- Change in consciousness
- Sensitivity of the eye to light
- Discharge, excessive tearing

- Confusion about identity, place, and time
- Red skin lesion often with a purple center

Consult your doctor if you face any of the side effects worsen.

Dosage of Ivermectin 12mg

Ivermectin 12mg medicine dose will different-different for the patient to patient. Because all the patient's body structures different-different so dose you have to follow the doctor's instructions. Also, the following information includes only the average doses of this medicine. Do not change your dose before talking with your doctor or unless your doctor tells you to do so.

Symptoms of overdose

- Confusion about identity, place, and time
- Decreased awareness or responsiveness
- Severe sleepiness

Instruction for Use

Whenever using Ivermectin 12mg tablet always take the advice of your doctor.

Ivermectin tablets should be taken as directed and with a glass of water. Don't chew or crush the medication. However, the medication needs to be crushed before being swallowed by children under the age of six.

Your doctor will determine the dosage and duration for you based on your age, body weight, and disease state.

Related Products

- [Ivermectin 6mg](#)
- Stromectol (Ivermectin 12 Mg)
- Ivermectin 3mg

Ivermectin 12mg Tablet FAQs:

Q: What is the Ivermectin 12mg tablet used for?

A: Ivermectin tablet is a widely used medication to treat various infections caused by parasites, including strongyloidiasis (a roundworm infection) and control onchocerciasis (a worm infection that affects skin and eyes).

Q: Is Ivermectin 12mg tablet safe during pregnancy?

A: No, It is not safe for use during pregnancy. Unless it is clearly advised by the doctor. Despite the Ivermectin drug's multiple health benefits. Pregnant women are typically excluded from ivermectin administration due to its potential effect on the foetus. Consult your doctor before taking an Ivermectin tablet. If you are pregnant or planning to get pregnant.

Q: Can [Ivermectin](#) tablets be used in children?

A: The safety and effectiveness of the Ivermectin tablet in children weighing less than 15 kg have not been evaluated. Your child's doctor may prescribe safer alternatives for your child to treat the specific infection. In the treatment of head lice, ivermectin topical lotion can be used on children 6 months of age and older.

Q: What precautions do I need to follow while taking the Ivermectin tablet?

A: Before taking an Ivermectin tablet. Inform your doctor if you are facing an eye infection, a weak immune system, or any allergies to any other medicines or any other substances such as foods, preservatives, or dyes.

Q: Can I drive or operate any machines after taking the Ivermectin tablet?

A: No, Do not drive or operate any machines if you feel dizziness, drowsiness, or feeling shaky or spinning sensation after taking the Ivermectin tablet. Contact your doctor before taking Ivermectin 12mg tablet.